BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:		(11) International Publication Number:	WO 97/38663
A61K	A2	(43) International Publication Date:	23 October 1997 (23.10.97)

(21) International Application Number: PCT/US97/02795

(22) International Filing Date: 21 February 1997 (21.02.97)

(30) Priority Data:

08/631,175

12 April 1996 (12.04.96)

US

(71) Applicant (for all designated States except US): FLEMING-TON PHARMACEUTICAL CORPORATION [US/US]; 43 Emery Avenue, Flemington, NJ 08822 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): DUGGER, Harry, A., III [US/US]; 548 Sargentville Road, Flemington, NJ 08822 (US).

(74) Agent: BEHR, Omri, M.; 325 Pierson Avenue, Edison, NJ 08837 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

Without international search report and to be republished upon receipt of that report.

(54) Title: BUCCAL, NON-POLAR SPRAY OR CAPSULE

(57) Abstract

A buccal aerosol spray or capsule using a non-polar solvent has now been developed which provides biologically active compounds for rapid absorption through the oral mucosa, resulting in fast onset of effect. The buccal aerosol spray of the invention comprises formulation (I): propellant 50-95 %, non-polar solvent 5-50 %, active compound 0.001-15 %, flavoring agent 0.05-5 %. The soft bite gelatin capsule of the invention comprises formulation (II): non-polar solvent 55-99.8 %, emulsifier 0-20 %, active compound 0.001-25 %, and flavoring agent 0.05-5.0 %.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	Codes used to identify	omico par	.,		_		
	A Thursday	ES	Spain	LS	Lesotho	SI	Slovenia
AL	Albania	Fi	Finland	LT	Lithuania	SK	Slovakia
AM	Armenia	FR	France	LU	Luxembourg	SN	Senegal
AT	Austria	GA	Gabon	LV	Latvia	SZ	Swaziland
AU	Australia	GB	United Kingdom	MC	Monaco	TD	Chad
ΑZ	Azerbaijan			MD	Republic of Moldova	TG	Togo
BA	Bosnia and Herzegovina	GE	Georgia Ghana	MG	Madagascar	TJ	Tajikistan
BB	Barbados	GH		MK	The former Yugoslav	TM	Turkmenistan
BB	Belgium	GN	Guinea	17124	Republic of Macedonia	TR	Turkey
BF	Burkina Faso	GR	Greece	ML	Mali	TT	Trinidad and Tobago
BG	Bulgaria	HU	Hungary	MN	Mongolia	UA	Ukraine
BJ	Benin	IE	Ireland	MR	Mauritania	UG	Uganda
BR	Brazil	IL	lsrael		Malawi	US	United States of America
BY	Belarus	IS	Iceland	MW		UZ	Uzbekistan
CA	Canada	IT	Italy	MX	Mexico	VN	Viet Nam
CF	Central African Republic	JP	Japan	NE	Niger	YU	Yugoslavia
CG	Congo	KB	Kenya	NL	Netherlands	zw	Zimbabwe
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	244	Zillozowo
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	u	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
	Batonia .	LR	Liberia	SG	Singapore		
EE	DECOME						

TITLE OF THE INVENTION BUCCAL, NON-POLAR SPRAY OR CAPSULE

BACKGROUND OF THE INVENTION

It is known that certain biologically active compounds are better 5 absorbed through the oral mucosa than through other routes of administration, such as through the stomach or intestine. However. formulations suitable for such administration by these latter routes present their own problems. For example, the biologically active compound must 10 be compatible with the other components of the composition such as propellants, solvents, etc. Many such formulations have been proposed. For example, U.S.P. 4,689,233, Dvorsky et al., describes a soft gelatin capsule for the administration of the anti-coronary drug nifedipine dissolved in a mixture of polyether alcohols. U.S.P. 4,755,389, Jones et al., 15 describes a hard gelatin chewable capsule containing nifedipine. A chewable gelatin capsule containing a solution or dispersion of a drug is described in U.S.P. 4,935,243, Borkan et al. U.S.P. 4,919,919, Aouda et al, and U.S.P. 5,370,862, Klokkers-Bethke, describe a nitroglycerin spray for administration to the oral mucosa comprising nitroglycerin, ethanol, and 20 other components. An orally administered pump spray is described by Aerosol compositions containing a Cholcha in U.S.P. 5,186,925. hydrocarbon propellant and a drug for administration to a mucosal surface are described in U.K. 2,082,457, Su, U.S.P. 3,155,574, Silson et al., U.S.P. 5,011,678, Wang et al., and by Parnell in U.S.P. 5,128,132. It 25 should be noted that these references discuss bioavailability of solutions by inhalation rather than through the membranes to which they are administered.

SUMMARY OF THE INVENTION

A buccal aerosol spray or soft bite gelatin capsule using a non-polar solvent has now been developed which provides biologically active com-

pounds for rapid absorption through the oral mucosa, resulting in fast onset of effect.

The buccal aerosol spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable non-polar solvent comprising in weight % of total composition: pharmaceutically acceptable propellant 50-95%, non-polar solvent 5-50%, active compound 0.0025-40%, suitably additionally comprising, by weight of total composition a flavoring agent 0.05-5%. Preferably the composition comprises: propellant 55-85%, non-polar solvent 15-45%, active compound 0.025-20%, flavoring agent 0.1-2.5%; most suitably propellant 60-80%, non-polar solvent 19-32%, active compound 0.125-12.5%, flavoring agent 0.1-3.0%.

The soft bite gelatin capsules of the present invention for transmucosal administration of a pharmacologically active compound, at least partially soluble in a pharmacologically acceptable non-polar solvent, having charged thereto a fill composition comprising in weight % of total composition: non-polar solvent 30-99.8%, emulsifier 0-20%, active compound 0.0003-32%, provided that said fill composition contains less than 10% of water, suitably additionally comprising, by weight of the composition: flavoring agent 0.05-60%. Preferably, the soft bite gelatin capsule comprises: non-polar solvent 40-99.8%, emulsifier 0-15%, active compound 0.004-26%, flavoring agent 0.1-55%; most suitably: non-polar solvent 40-99.5%, emulsifier 0-10%, active compound 0.015-24.0%, flavoring agent 0.1-50%.

It is an object of the invention to coat the mucosal membranes either with extremely fine droplets of spray containing the active compounds or a solution or paste thereof from bite capsules.

WO 97/38663 PCT/US97/02795

It is also an object of the invention to administer to a mammalian in need of same preferably man, a predetermined amount of a biologically active compound by this method or from a soft gelatin bite capsule.

A further object is a sealed aerosol spray container containing a composition of the spray formulation, and a metered valve suitable for releasing from said container a predetermined amount of said composition.

As the propellant evaporates after activation of the aerosol valve, a 10 mist of fine droplets is formed which contains solvent and active compound.

The propellant is a non-Freon material, preferably a C₃₋₈ hydrocarbon of a linear or branched configuration. The propellant should be substantially non-aqueous. The propellant produces a pressure in the aerosol container such that under expected normal usage it will produce sufficient pressure to expel the solvent from the container when the valve is activated but not excessive pressure such as to damage the container or valve seals.

The solvent is a non-polar hydrocarbon, preferably a C₇₋₁₈ hydrocarbon 20 of a linear or branched configuration, its alcohols, fatty acid esters, and triglycerides, such as miglyol. The solvent must dissolve the active compound and be miscible with the propellant, i.e., solvent and propellant must form a single phase at 0-40°C at a pressure range of 1-3 atm.

A further object is a soft gelatin bite capsule containing a composition of as set forth above. The formulation may be in the form of a viscous solution or paste containing the active compounds. Although solutions are preferred, paste fills may also be used where the active compound is not soluble or only partially soluble in the solvent of choice. Where water is 30 used to form part of the paste composition, it should not exceed 10% thereof. (All percentages herein are by weight unless otherwise indicated.)

The non-polar solvent is chosen such that it is compatible with the gelatin shell and the active compound. The solvent preferably dissolves the active compound. However, other components wherein the active compound is not soluble or only slightly soluble may be used and will form a paste fill.

Soft gelatin capsules are well known in the art. See, for example, U.S.P. 4,935,243, Borkan et al., which is incorporated herein by reference for its teaching of such capsules. The capsules of the present invention are 10 intended to be bitten into to release the low viscosity solution or paste therein, which will then coat the buccal mucosa with the active compounds. Typical capsules, which are swallowed whole or bitten and then swallowed, deliver the active compounds the stomach, which results in significant lag time before maximum blood levels can be achieved or subject the compound 15 to a large first pass effect. Because of the enhanced absorption of the compounds through the oral mucosa and no chance of a first pass effect, use of the bite capsules of the invention will eliminate much of the lag time, resulting in hastened onset of biological effect. The shell of a soft gelatin capsule of the invention may comprise, for example: gelatine 50-75%, glycerine 20-30%, colorants 0.5-1.5%, water 5-10%, and sorbitol 2-10%.

The spray compositions of the invention are intended to be administered from a sealed, pressurized container. Unlike a pump spray, which allows the entry of air into the container after every activation, the aerosol container of the invention is sealed at the time of manufacture. The contents of the container are released by activation of a metered valve, will does not allow entry of atmospheric gasses with each activation. Such containers are commercially available.

WO 97/38663 PCT/US97/02795 5

BRIEF DESCRIPTION OF THE DRAWING

The figure is a schematic diagram showing routes of absorption and processing of pharmacologically active substances in a mammalian system.

5 <u>DESCRIPTION OF THE PREFERRED EMBODIMENTS</u>

The preferred active compounds of the present invention are nicotine, clemastine, testosterone, estradiol, progesterone, flüoxetine, and piroxicam in their nonionized form or as the free base of the pharmaceutically acceptable salts thereof (provided, for the aerosol or spray compositions, they are soluble in the spray solvent), as well as, where appropriate the esters or triglycerides thereof. These compounds are soluble in the non-polar solvents of the invention at useful concentrations or can be prepared as pastes at useful concentrations. These concentrations may be less than the standard accepted dose for these compounds since there is enhanced absorption of the compounds through the oral mucosa. This aspect of the invention is especially important when there is a large (40-99.99%) First pass effect.

As propellants for the sprays, propane, N-butane, iso-butane, 20 N-pentane, iso-pentane, and neo-pentane, and mixtures thereof may be used. N-butane and iso-butane, as single gases, are the preferred propellants. It is permissible for the propellant to have a water content of no more than 0.2%, typically 0.1-0.2%. (All percentages herein are by weight unless otherwise indicated.) It is also preferable that the propellant 25 be synthetically produced to minimize the presence of contaminants which are harmful to the active compounds. These contaminants include oxidizing agents, reducing agents, Lewis acids or bases, and water. The concentration of each of these should be less than 0.1%, except that water may be as high as 0.2%.

Suitable solvents for the capsules include the spray solvents listed above, as well as [C₂-C₂₄] fatty acid C₂-C₆ esters, including the triglycerides. Similarly, water is not a suitable solvent component in the spray compositions. When the capsule fill is a paste, other liquid components may be used instead of the above low molecular weight solvents. These include soya oil, corn oil, other vegetable oils, and C₇₋₁₈ hydrocarbons of a linear or branched configuration, and their alcohols and their fatty acid esters and triglycerides.

10 It is expected that some glycerin and water used to make the gelatin shell will migrate from the shell to the fill during the curing of the shell. Likewise, there may be some migration of components from the fill to the shell during curing and even throughout the shelf-life of the capsule. Therefore, the values given herein are for the compositions as prepared, it being within the scope of the invention that minor variations will occur.

The preferred flavoring agents are synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, sweeteners (sugars, aspartame, saccharin, etc.), and combinations thereof.

20

The active substances include the active compounds selected from the group consisting of alkaloids, anti histamines, steroid hormones, non-steroidal anti-inflammatories, analgesics and anti-depressants, benzo-diazepines, such as tamezepam.

25

Clemastine hydrogen fumarate is a known (Tavist[®], Sandoz) antihistamine. Both the spray and capsule of the invention advantageously coat the oral mucosa with an immediately available dose of clemastine which can be rapidly absorbed. This is highly desirable, as during an acute asthma 30 attack. Nicotine is a component of tobacco products which is considered addictive. Smokers wishing to stop smoking have a dual problem. First, is the addictive properties of nicotine itself. Second, is that the habit is associated with smoking activities, i.e., puffing, inhaling, etc. Both the spray and capsule of the invention dissociate these two problems. By presenting nicotine in a form which can be readily absorbed, the spray and capsule allow the smoker to temporarily continue nicotine use but terminate smoking. Once the habit of smoking is stopped, the former smoker can then be weaned off nicotine use, as by less frequent use and/or by use of a lower concentration spray or capsule. Advantageously, during this regimen, the user is exposed to none of the carcinogens present in tobacco smoke.

Testosterone is a hormone produced by gonadal cells. Testosterone, 15 especially the esters thereof (e.g., acetate, propionate, enanthate, and cypionate), is used in the treatment of hypogonadism.

Estradiol is an estrogen steroid secreted from the ovaries. Estradiol, especially the esters thereof (e.g., diacetate, and benzoate), is used as 20 estrogen replacement therapy, especially in post-menopausal women.

Progesterone is a hormone produced by the corpus luteum. Fluoxetine is an antidepressant also known as Prozac. Piroxicam is a known (Feldene®, Pfizer) anti-inflammatory.

25

The formulations of the present invention comprise an active compound or a pharmaceutically acceptable salt thereof. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic acids or bases including organic and inorganic acids or bases.

When an active compound of the present invention is acidic, salts may be prepared from pharmaceutically acceptable non-toxic bases. Salts derived from all stable forms of inorganic bases include aluminum, ammonium, calcium, copper, iron, lithium, magnesium, manganese, 5 potassium, sodium, zinc, etc. Particularly preferred are the ammonium. calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion-10 exchange resins such as arginine, betaine, caffeine, choline, N,N'dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, isopropylamine, lysine, methylglucosamine, morpholine, piperazine, piperidine, polyamine resins, 15 procaine, purine, theobromine, triethylamine, trimethylamine, tripropylamine, etc.

When an active compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic, etc. Particularly preferred are citric, hydrobromic, maleic, phosphoric, sulfuric, and tartaric acids.

In the discussion of methods of treatment herein, reference to the active compounds is meant to also include the pharmaceutically acceptable salts thereof. While certain formulations are set forth herein, the actual armounts to be administered to the mammal or man in need of same are to be determined by the treating physician.

The invention is further defined by reference to the following examples, which are intended to be illustrative and not limiting.

EXAMPLE 1

5 Clemastine Spray

A spray of the invention comprises the following formulation:

		Amount	Preferred Amount	Most-Preferred Amount
	Propellant	50-95%	55-85%	65-80%
	Non-polar solvent	5-50%	15-45%	20-35%
10	Clemastine Base	0.12-10%	0.25-6.25%	0.25-5%
	Flavoring agent	0.05-3%	0.1-2.5%	0.2-2%

It is particularly preferred to formulate the spray delivering 1.0mg/activation formulation:

15		<u>Amount</u>
	Butane	67%
	Miglyol	29.5%
	Clemastine Base	3.1%
	Peppermint	0.4%

For delivery of 0.5mg active substance, Clemastine Base 1.6%

EXAMPLE 2

Nicotine Spray

A spray of the invention comprises the following formulation:

25		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Propellant	50-95%	55-85%	65-80%
	Non-polar solvent	5-50%	15-45%	20-35%
	Nicotine	0.125-2.5%	0.25-1.75%	0.25-1.25%
	Flavoring agent	0.05-3%	0.1-2.5%	0.1-2.5%

It is particularly preferred to formulate the spray delivering 0.5mg/activation:

<u>Amount</u>

Butane

67%

5 Miglyol

31.25%

Nicotine

1.5% for 0.5mg.

Peppermint

0.25%

For delivery of 0.4mg active substance, Nicotine 1.25%.

10 EXAMPLE 3

Testosterone Spray

A spray of the invention comprises the following formulation:

		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Propellant	50-95%	55-85%	65-80%
15	Non-polar solvent	5-50%	15-45%	20-35%
	Testosterone*	0.125-25%	2.5-20%	5-12.5%
	Flavoring agent	0.05-3%	0.1-2.5%	0.1-2.5%

^{*}or esters thereof, preferably, the acetate, propionate, and enenthate esters

20 It is particularly preferred to formulate the spray delivering 4mg/ activation:

		<u>Amount</u>
	Butane	67%
	Miglyol	20.25%
25	Testosterone	12.5%
	Peppermint	0.25%

EXAMPLE 4

Estradiol Spray

A spray of the invention comprises the following formulation:

		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
5	Propellant	50-75%	55-85%	75-99%
	Non-polar solvent	25-50%	15-45%	20-35%
	Estradiol*	0.0025-2.59	% 0.025-1. 5 %	0.125-1.0%
	Flavoring agent	0.05-3%	0.1-2.5%	0.1-2.5%

^{*}or esters thereof, preferably, the diacetate and benzoate esters

10

It is particularly preferred to formulate the spray:

		<u>Amount</u>
	Butane	67%
	Miglyol	32.4%
15	Estradiol	0.31%
	Peppermint	0.25%

EXAMPLE 5

Progesterone Spray

A spray of the invention comprises the following formulation:

		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Propellant	50-75%	55-85%	65-80%
	Non-polar solvent	25-50%	15-45%	20-35%
	Progesterone	0.0025-2.59	0.025-2.0%	0.125-0.5%
25	Flavoring agent	0.05-3%	0.1-2.5%	0.1-2.5%

It is particularly preferred to formulate the spray delivering 0.32mg/ activation:

		<u>Amount</u>
	Butane	67%
5	Miglyol	31.2%
	Progesterone	1.0%
	Spearmint	0.5%

EXAMPLE 6

10 Clemastine Bite Capsule

A paste containing bite capsule of the invention comprises the following fill formulation:

		Amount F	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	75-99%	90-98%	85-99.5%
15	Emulsifier	0-20%	0-15%	0-10%
	Clemastine fumara	te' 0.0003-1.8	35% 0.003-0.74	% 0.018-0.185%
	Flavoring agent	0.05-5%	0.1-2.5%	0.1-2.5%
	'as paste			

capsule:

	It is particula	arly preferred to formulate the fill for the 1.34mg ca
20		Amount
	Soya Oil	91.4%
	Lecithin	5.88%
	Clemastine fumara	ate 0.50% for 1.34mg.
	Orange Aroma	1.04%
25	Citron Oil	0.81%
	Na-Saccharine	0.37%

For delivery of 0.68m active substance, Clemastine fumarate 0.25%.

EXAMPLE 7

Testosterone Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

5		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	55-99%	66-97%	85-99.5%
	Emulsifier/wetting	0-20%	0-15%	0-10%
	agents			
	Testosterone*	0.01-3.7%	0.37-2.96%	0.7-1.85%
10	Flavoring agent	0.05-5%	0.1-2.5%	0.1-2.5%

^{*}or esters thereof, preferably, the acetate, propionate, and enenthate esters

It is particularly preferred to formulate the fill for 5mg capsule:

		<u>Amount</u>
15	Miglyol	90.1%
	Lecithin	5.88%
	Testosterone	1.8%
	Orange Aroma	1.04%
	Citron Oil	0.81%
20	Na-Saccharine	0.37%

EXAMPLE 8

Estradiol Bite Capsule

A bite capsule of the invention comprises the following fill 25 formulation:

		Amount	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	65-99%	85-98%	85-99.5%
30	Emulsifier/wetting agents	0-20%	0-15%	0-10%
	Estradiol*	0.0003-1.85	% 0.003-0.749	% 0.018-0.185%
	Flavoring agent	0.05-5%	0.1-2.5%	0.1-2.5%

14

*or esters thereof, preferably, the diacetate and benzoate esters

It is particularly preferred to formulate the fill for 0.5mg capsule:

		<u>Amount</u>
5	Miglyol	91.71%
	Lecithin	5.88%
	Estradiol	0.18 for 0.5mg capsules
	Orange Aroma	1.04%
	Citron Oil	0.81%
10	Na Saccharine	0.37%

EXAMPLE 9

Progesterone Bite Capsule

A bite capsule of the invention comprises the following fill 15 formulation:

		Amount E	referred Amount	Most-Preferred Amount
	Non-polar solvent	55-99%	75-98%	85-99.5%
	Emulsifier/wetting	0-20%	0-15%	0-10%
	agents			
20	Progesterone	0.0003-3.7%	0.0003-2.79	% 0.74-1.85%
	Flavoring agent	0.05-5%	0.1-2.5%	0.1-2.5%

It is particularly preferred to formulate the fill for a 3mg capsule:

		<u>Amount</u>
25	Miglyol	91.83%
	Lecithin	5.88%
	Progesterone	1.11%
	Oil of Citron	0.81%
	Na-Saccharine	0.37%

PCT/US97/02795 WO 97/38663

15

EXAMPLE 10

Fluoxetine Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

5		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	75-99.8%	75-99.8%	85-99.5%
	Emulsifier/wetting	0-20%	0-15%	0-10%
	agents			
	Fluoxetine* HCI**	0.018-9.259	% 0.37-3.7%	0.74-3.7%**
10	Flavoring agent	0.05-5%	0.1-3%	0.5-2.5%
	*If used as the fre	e base, it will	be a solution, at le	east in part

It is particularly preferred to formulate the fill for a 5mg capsule:

15		Amount
	Soya Oil	91.07%
	Lecithin	5.88%
	Fluoxetine HCI	1.85%
	Peppermint	1.2%

20

EXAMPLE 11

Piroxicam Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

25		Amount	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	75-99.8%	75-99.8%	85-99.5%
	Emulsifier/wetting	0-20%	0-15%	0-10%
	agents			
	Piroxicam	0.018-9.25%	6 0.37-3.7%	0.74-3.7%
30	Flavoring agent	0.05-5%	0.1-3%	0.5-2.5%

^{**} As a paste.

It is particularly preferred to formulate the fill for a 5mg capsule:

		Amount
	Soya Oil	91.07%
	Lecithin	5.88%
5	Piroxicam	1.85%
	Oil of Citron	0.81%
	Na Saccharine	0.37%

EXAMPLE 12

10 Nicotine 0.5mg Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

		<u>Amount</u>	Preferred Amount	Most-Preferred Amount
	Non-polar solvent	30-99.8%	35-99.8%	40-99.5%
15	Emulsifier/wetting	0-20%	0-15%	0-10%
	agents			
	Nicotine*	0.018-0.749	% 0.037-0.37	% 0.37-0.20%
	Flavoring agent	0.05-60%	0.1-55%	0.5-50%
	' or on pigoting cul	fato		

or as nicotine sulfate

20 It is particularly preferred to formulate the fill for a 0.5mg capsule:

		<u>Amount</u>
	Soya Oil	46.67%
	Chocolate powder	46.67%
	Lecithin	6.105%
25	Peppermint Oil	0.185%
	Na Saccharine	0.185%
	Nicotine	0.185%

17

EXAMPLE 13

Clemastine Fumarate with Phenylpropanolamine Hydrochloride Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

5		<u>Amount</u>	<u>Prefer</u>	red Amount	Most-	Preferred Amo	ount
	Non-Polar solvent	65-99%		50-98%		70-98%	
10	Clemastine fumara	te* 0-01-1.8	5%	0.03-0.74%		0.018-0.185	5%
	Phenylpropanolami HCI*	ne 1-30%		1.5-20%		1.8-10%	
	Flavoring agent	0.05-5%		0.1-2.5%		0.1-2.5%	
	Emulsifier/Wetting Agent	0-20%		0-15%		0-10%	

15 *Present as paste.

It is particularly preferred to formulate the composition fill for a clemastine fumarate 1.34mg/25mg phenylpropanolamine hydrochloride bite 20 capsule:

		<u>Amount</u>
	Non-Polar Solvent:	
	Soya	82.20%
	Lecithin	5.88%
25	Clemastine fumarate	0.50%
	Phenylpropanolamine HC	9.20%
	Orange aroma	1.04%
	Oil of Citrus	0.81%
	Na Saccharine	0.37%

30

EXAMPLE 14

Glemastine Fumarate/Pseudoephedrine Hydrochloride Bite Capsule

A bite capsule of the invention comprises the following fill formulation:

PCT/US97/02795

18

		<u>Amount</u>		red Amount	Most-Preferred Amount	
5	Non-Polar solvent	50-95%		55-90%	60-85%	
	Emulsifier/wetting agent	0-20%		0-15%	0-10%	
	Clemastine fumara	ite* 0.01-1.8	5%	0.03-0.74%	0.05-0.185%	
	Pseudoephedrine I	HCI* 3-30%		5-25%	10-23%	
	Flavoring agent	0.05-5%		0.1-2.5%	0.1-2.5%	
	*Present as paste	•				

10

It is particularly preferred to formulate the composition fill for a 1.34mg clemastine fumarate/60mg pseudoephedrine HCl capsule:

<u>Amount</u>

Non-Polar Solvent:

15	Soya	69.18%
	Lecithin	5.88%
	Clemastine fumarate	0.50%
	Phenylpropanolamine I	HCI 22.22%
	Orange aroma	1.04%
20	Oil of Citrus	0.81%
	Na Saccharine	0.37%

WO 97/38663 PCT/US97/02795

19

WHAT IS CLAIMED IS:

- A buccal aerosol spray composition for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable non-polar solvent comprising in weight % of total composition: pharmaceutically acceptable propellant 50-95%, non-polar solvent 5-50%, active compound 0.0025-40%.
- 2. The composition of claim 1 additionally comprising, by weight 10 of total composition: flavoring agent 0.05-5%.
 - 3. The composition of claim 1 comprising: propellant 55-85%, non-polar solvent 15-45%, active compound 0.025-20%, flavoring agent 0.1-2.5%.

- 4. The composition of claim 1 comprising: propellant 60-80%, non-polar solvent 19-32%, active compound 0.125-12.5%, flavoring agent 1-2%.
- 5. The composition of Claim 1 wherein the propellant is a C_{3-8} hydrocarbon of a linear or branched configuration.
- 6. The composition of Claim 1 wherein the propellant is propane, N-butane, iso-butane, N-pentane, iso-pentane, or neo-pentane, and mixtures thereof.
- 7. The composition of Claim 1 wherein the propellant is N-butane or iso-butane and has a water content of no more than 0.2% and oxidizing agents, reducing agents, and Lewis acids or bases content in a concentration of less than 0.1%

8. The composition of Claim 1 wherein the solvent is a selected from the group consisting of (C_2-C_{24}) fatty acid (C_2-C_6) esters, C_7-C_{18} hydrocarbons of a linear or branched configuration, the alcohols thereof, and C_2-C_6 alkanoyl esters, and triglycerides of said alcohols.

- 9. The composition of Claim 1 wherein the solvent is miglyol.
- 10. The composition of Claim 1 wherein the active compound is selected from the group consisting of alkaloids, anti-histamines, steroid
 10 hormones, non-steroidal anti-inflammatories, analgesics, benzediazepines and anti-depressants.
- 11. The composition of Claim 10 wherein the active compound is selected from the group consisting of nicotine, clemastine, testosterone,
 15 estradiol, progesterone, tamezepam, fluoxetine, and piroxicam in their nonionized form or as the free base of the pharmaceutically acceptable salts thereof.
- 12. The composition of Claim 2 wherein the flavoring agents are selected from the group consisting of synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, sweeteners and combinations thereof.
- 13. The composition of Claim 1 of the formulation: propellant25 50-95%, non-polar solvent 5-50%, clemastine base 0.12-10%, flavoring agent 0.05-3%.
- 14. The composition of Claim 1 of the formulation: propellant 50-95%, non-polar solvent 5-50%, nicotine 0.125-2.5%, flavoring agent 30 0.05-3%.

WO 97/38663 PCT/US97/02795

21

15. The composition of Claim 1 of the formulation: propellant 50-95%, non-polar solvent 5-50%, a member selected from the group

consisting of testosterone and the pharmaceutically acceptable esters

thereof 0.125-25%, flavoring agent 0.05-3%.

5

16. The composition of Claim 1 of the formulation: propellant 50-75%, non-polar solvent 25-50%, a member selected from the group consisting of estradiol and the pharmaceutically acceptable esters thereof 0.0025-2.5%, flavoring agent 0.05-3%.

10

17. The composition of Claim 1 of the formulation: propellant 50-75%, non-polar solvent 25-50%, a member selected from the group consisting of progesterone and the pharmaceutically acceptable esters thereof 0.025-2.5%, flavoring agent 0.05-3%.

- 18. A method of administering a pharmacologically active compound to a mammal in needed of same, by spraying the oral mucosa of said mammal with a composition of claim 1.
- 20 19. The method of claim 18 wherein the amount of spray administered is predetermined.
- 20. A sealed aerosol spray container containing a composition of claim 1 and a metered valve suitable for releasing from said container a
 25 predetermined amount of said composition.
- 21. A soft bite gelatin capsule for transmucosal administration of a pharmacologically active compound, at least partially soluble in a pharmacologically acceptable non-polar solvent, having charged thereto a 30 fill composition comprising in weight % of total fill composition: non-polar solvent 30-99.8%, emulsifier 0-20%, active compound 0.0003-32%,

5

provided that said composition contains less than 10% of water.

- 22. The capsule of claim 21 additionally comprising, by weight of the fill composition: flavoring agent 0.05-60%.
- 23. The soft bite gelatin capsule of Claim 22 comprising: non-polar solvent 40-99.8%, emulsifier 0-15%, active compound 0.004-26%, flavoring agent 0.1-5%.
- 10 24. The soft bite gelatin capsule of Claim 22 as fill composition comprising: non-polar solvent 40-99.5%, emulsifier 0-10%, active compound 0.015- 24.0%, flavoring agent 0.1-50%.
- 25. The capsule of Claim 21 wherein the solvent is selected from 15 the group consisting of (C_2-C_{24}) fatty acid (C_2-C_8) esters, C_7-C_{18} hydrocarbons of a linear or branched configuration, the alcohols thereof, and C_2-C_8 alkanoyl esters, and triglycerides of said alcohols.
- 26. The capsule of Claim 21 wherein the solvent is selected from 20 the group consisting of alkaloids, anti histamines, steroid hormones, nonsteroidal anti-inflammatories, analgesics, benzediazepines and antidepressants.
- 27. The capsule of Claim 21 wherein the active compound is 25 nicotine, clemastine, testosterone, estradiol, progesterone, fluoxetine, temazepam, or piroxicam in their nonionized form or as the free base of the pharmaceutically acceptable salts thereof.
- 28. The capsule of Claim 21 wherein the flavoring agents are 30 synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, or sweeteners and combinations thereof.


WO 97/38663 PCT/US97/02795

- 29. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 75-99%, emulsifier/wetting 0-20%, clemastine fumarate 0.01-4%, flavoring agent 0.05-5%.
- 5 30. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 55-99%, emulsifier 0-20%, A member selected from the group consisting of testosterone and the esters thereof 0.01-4%, flavoring agent 0.05-5%.
- 10 31. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 65-99%, emulsifier/wetting 0-20%, estradiol 0.00031- 2%, flavoring agent 0.05-5%.
- 32. The capsule of Claim 27 comprising fill composition of the 15 formulation: non-polar solvent 55-99%, emulsifier/wetting 0-20%, progesterone 0.0003-4%, flavoring agent 0.05-5%.
- 33. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 75-99.8%, emulsifier 0-20%, fluoxetine HCl20 0.02-9.5%, flavoring agent 0.05-5%.
 - 34. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 75-99.8%, emulsifier 0-20%, piroxicam 0.02-9.5%, flavoring agent 0.05-5%.

- 34. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 30-99.8%, emulsifier 0-20%, nicotine 0.018 0.74%, flavoring agent 0.05-60%.
- 35. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 65-99.8%, emulsifier 0-20%, phenyl

propanolamine 1-30%, clemastine 0.01 - 1.85% flavoring agent 0.05-5%.

36. The capsule of Claim 27 comprising fill composition of the formulation: non-polar solvent 50-95%, emulsifier 0-20%, pseudoephedrine
5 3-30%, clemastine 0.01 - 1.85% flavoring agent 0.05-5%.



K(e) = K(m) + K(b) + K(u)

FIGURE